

UPSC Civil Services Main 1995 - Mathematics

Linear Algebra

Sunder Lal

Retired Professor of Mathematics

Panjab University

Chandigarh

June 14, 2007

1 Linear Algebra

Question 1(a) Let $\mathbf{T}(x_1, x_2, x_3) = (3x_1 + x_3, -2x_1 + x_2, -x_1 + 2x_2 + 4x_3)$ be a linear transformation on \mathbb{R}^3 . What is the matrix of \mathbf{T} w.r.t. the standard basis? What is a basis of the range space of \mathbf{T} ? What is a basis of the null space of \mathbf{T} ?

Solution.

$$\begin{aligned}\mathbf{T}(\mathbf{e}_1) &= \mathbf{T}(1, 0, 0) = (3, -2, -1) = 3\mathbf{e}_1 - 2\mathbf{e}_2 - \mathbf{e}_3 \\ \mathbf{T}(\mathbf{e}_2) &= \mathbf{T}(0, 1, 0) = (0, 1, 2) = \mathbf{e}_2 + 2\mathbf{e}_3 \\ \mathbf{T}(\mathbf{e}_3) &= \mathbf{T}(0, 0, 1) = (1, 0, 4) = \mathbf{e}_1 + 4\mathbf{e}_3 \\ \mathbf{T} \iff \mathbf{A} &= \begin{pmatrix} 3 & 0 & 1 \\ -2 & 1 & 0 \\ -1 & 2 & 4 \end{pmatrix}\end{aligned}$$

Clearly $\mathbf{T}(\mathbf{e}_2), \mathbf{T}(\mathbf{e}_3)$ are linearly independent. If $(3, -2, -1) = \alpha(0, 1, 2) + \beta(1, 0, 4)$, then $\beta = 3, \alpha = -2$, but $2\alpha + 4\beta \neq -1$, so $\mathbf{T}(\mathbf{e}_1), \mathbf{T}(\mathbf{e}_2), \mathbf{T}(\mathbf{e}_3)$ are linearly independent. Thus $(3, -2, -1), (0, 1, 2), (1, 0, 4)$ is a basis of the range space of \mathbf{T} .

Note that $\mathbf{T}(x_1, x_2, x_3) = \mathbf{0} \iff x_1 = x_2 = x_3 = 0$, so the null space of \mathbf{T} is $\{\mathbf{0}\}$, and the empty set is a basis. Note that the matrix of \mathbf{T} is nonsingular, so $\mathbf{T}(\mathbf{e}_1), \mathbf{T}(\mathbf{e}_2), \mathbf{T}(\mathbf{e}_3)$ are linearly independent. ■

Question 1(b) Let \mathbf{A} be a square matrix of order n . Prove that $\mathbf{Ax} = \mathbf{b}$ has a solution $\iff \mathbf{b} \in \mathbb{R}^n$ is orthogonal to all solutions \mathbf{y} of the system $\mathbf{A}'\mathbf{y} = \mathbf{0}$.

Solution. If \mathbf{x} is a solution of $\mathbf{Ax} = \mathbf{b}$ and \mathbf{y} is a solution of $\mathbf{A}'\mathbf{y} = \mathbf{0}$, then $\mathbf{b}'\mathbf{y} = \mathbf{x}'\mathbf{A}'\mathbf{y} = 0$, thus \mathbf{b} is orthogonal to \mathbf{y} .

Conversely, suppose $\mathbf{b}'\mathbf{y} = 0$ for all $\mathbf{y} \in \mathbb{R}^n$ which is a solution of $\mathbf{A}'\mathbf{y} = \mathbf{0}$. Let $\mathcal{W} = \mathbf{A}(\mathbb{R}^n)$ = the range space of \mathbf{A} , and \mathcal{W}^\perp its orthogonal complement. If $\mathbf{A}'\mathbf{y} = \mathbf{0}$ then $\mathbf{x}'\mathbf{A}'\mathbf{y} = 0 \Rightarrow (\mathbf{Ax})'\mathbf{y} = 0$ for every $\mathbf{x} \in \mathbb{R}^n \Rightarrow \mathbf{y} \in \mathcal{W}^\perp$. Conversely $\mathbf{y} \in \mathcal{W}^\perp \Rightarrow \forall \mathbf{x} \in \mathbb{R}^n. (\mathbf{Ax})'\mathbf{y} = 0 \Rightarrow \mathbf{x}'\mathbf{A}'\mathbf{y} = 0 \Rightarrow \mathbf{A}'\mathbf{y} = \mathbf{0}$. Thus $\mathcal{W}^\perp = \{\mathbf{y} \mid \mathbf{A}'\mathbf{y} = \mathbf{0}\}$. Now $\mathbf{b}'\mathbf{y} = 0$ for all $\mathbf{y} \in \mathcal{W}^\perp$, so $\mathbf{b} \in \mathcal{W} \Rightarrow \mathbf{b} = \mathbf{Ax}$ for some $\mathbf{x} \in \mathbb{R}^n \Rightarrow \mathbf{Ax} = \mathbf{b}$ is solvable. ■

Question 1(c) Define a similar matrix and prove that two similar matrices have the same characteristic equation. Write down a matrix having 1, 2, 3 as eigenvalues. Is such a matrix unique?

Solution. Two matrices \mathbf{A}, \mathbf{B} are said to be similar if there exists a matrix \mathbf{P} such that $\mathbf{B} = \mathbf{P}^{-1}\mathbf{A}\mathbf{P}$. If \mathbf{A}, \mathbf{B} are similar, say $\mathbf{B} = \mathbf{P}^{-1}\mathbf{A}\mathbf{P}$, then characteristic polynomial of \mathbf{B} is $|\lambda\mathbf{I} - \mathbf{B}| = |\lambda\mathbf{I} - \mathbf{P}^{-1}\mathbf{A}\mathbf{P}| = |\mathbf{P}^{-1}\lambda\mathbf{I}\mathbf{P} - \mathbf{P}^{-1}\mathbf{A}\mathbf{P}| = |\mathbf{P}^{-1}||\lambda\mathbf{I} - \mathbf{A}||\mathbf{P}| = |\lambda\mathbf{I} - \mathbf{A}|$. (Note that $|\mathbf{X}||\mathbf{Y}| = |\mathbf{XY}|$.) Thus the characteristic polynomial of \mathbf{B} is the same as that of \mathbf{A} .

Clearly the matrix $\mathbf{A} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{pmatrix}$ has eigenvalues 1, 2, 3. Such a matrix is not unique, for example $\mathbf{B} = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{pmatrix}$ has the same eigenvalues, but $\mathbf{B} \neq \mathbf{A}$. ■

Question 2(a) Show that

$$\mathbf{A} = \begin{pmatrix} 5 & -6 & -6 \\ -1 & 4 & 2 \\ 3 & -6 & -4 \end{pmatrix}$$

is diagonalizable and hence determine \mathbf{A}^5 .

Solution.

$$\begin{aligned} & |\mathbf{A} - \lambda\mathbf{I}| = 0 \\ \Rightarrow & \begin{vmatrix} 5 - \lambda & -6 & -6 \\ -1 & 4 - \lambda & 2 \\ 3 & -6 & -4 - \lambda \end{vmatrix} = 0 \\ \Rightarrow & (5 - \lambda)[(4 - \lambda)(-4 - \lambda) + 12] + 6[4 + \lambda - 6] - 6[6 - 3(4 - \lambda)] = 0 \\ \Rightarrow & (5 - \lambda)[\lambda^2 - 4] + 6[\lambda - 2 - 3\lambda + 6] = 0 \\ \Rightarrow & -\lambda^3 + 5\lambda^2 + 4\lambda - 20 - 12\lambda + 24 = 0 \\ \Rightarrow & \lambda^3 - 5\lambda^2 + 8\lambda - 4 = 0 \end{aligned}$$

Thus $\lambda = 1, 2, 2$.

If (x_1, x_2, x_3) is an eigenvector for $\lambda = 1$, then

$$\begin{aligned} & \begin{pmatrix} 4 & -6 & -6 \\ -1 & 3 & 2 \\ 3 & -6 & -5 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \mathbf{0} \\ \Rightarrow & 4x_1 - 6x_2 - 6x_3 = 0 \\ & -x_1 + 3x_2 + 2x_3 = 0 \\ & 3x_1 - 6x_2 - 5x_3 = 0 \end{aligned}$$

Thus $x_1 = x_3, x_3 = -3x_2$, so $(-3, 1, -3)$ is an eigenvector for $\lambda = 1$.

If (x_1, x_2, x_3) is an eigenvector for $\lambda = 2$, then

$$\begin{aligned} \begin{pmatrix} 3 & -6 & -6 \\ -1 & 2 & 2 \\ 3 & -6 & -6 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} &= \mathbf{0} \\ \Rightarrow 3x_1 - 6x_2 - 6x_3 &= 0 \\ -x_1 + 2x_2 + 2x_3 &= 0 \\ 3x_1 - 6x_2 - 6x_3 &= 0 \end{aligned}$$

Thus $x_1 - 2x_2 - 2x_3 = 0$, so taking $x_1 = 0, x_2 = 1$, $(0, 1, -1)$ is an eigenvector for $\lambda = 2$. Taking $x_1 = 4, x_2 = 1$, $(4, 1, 1)$ is another eigenvector for $\lambda = 2$, and these two are linearly independent.

Let $\mathbf{P} = \begin{pmatrix} -3 & 0 & 4 \\ 1 & 1 & 1 \\ -3 & -1 & 1 \end{pmatrix}$. A simple calculation shows that $\mathbf{P}^{-1} = \frac{1}{2} \begin{pmatrix} 2 & -4 & -4 \\ -4 & 9 & 7 \\ 2 & -3 & -3 \end{pmatrix}$.

Clearly $\mathbf{P}^{-1}\mathbf{A}\mathbf{P} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{pmatrix}$.

Now $\mathbf{P}^{-1}\mathbf{A}^5\mathbf{P} = (\mathbf{P}^{-1}\mathbf{A}\mathbf{P})^5 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 32 & 0 \\ 0 & 0 & 32 \end{pmatrix}$.

$$\begin{aligned} \mathbf{A}^5 &= \mathbf{P} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 32 & 0 \\ 0 & 0 & 32 \end{pmatrix} \mathbf{P}^{-1} \\ &= \frac{1}{2} \begin{pmatrix} -3 & 0 & 4 \\ 1 & 1 & 1 \\ -3 & -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 32 & 0 \\ 0 & 0 & 32 \end{pmatrix} \begin{pmatrix} 2 & -4 & -4 \\ -4 & 9 & 7 \\ 2 & -3 & -3 \end{pmatrix} \\ &= \frac{1}{2} \begin{pmatrix} -3 & 0 & 128 \\ 1 & 32 & 32 \\ -3 & -32 & 32 \end{pmatrix} \begin{pmatrix} 2 & -4 & -4 \\ -4 & 9 & 7 \\ 2 & -3 & -3 \end{pmatrix} \\ &= \begin{pmatrix} 125 & -186 & -186 \\ -31 & 94 & 62 \\ 93 & -186 & -154 \end{pmatrix} \end{aligned}$$

Note: Another way of computing \mathbf{A}^5 is given below. This uses the characteristic polynomial of \mathbf{A} : $\mathbf{A}^3 = 5\mathbf{A}^2 - 8\mathbf{A} + 4\mathbf{I}$ and not the diagonal form, so it will *not* be permissible here.

$$\begin{aligned}
\mathbf{A}^5 &= \mathbf{A}^2(5\mathbf{A}^2 - 8\mathbf{A} + 4\mathbf{I}) \\
&= 5\mathbf{A}(5\mathbf{A}^2 - 8\mathbf{A} + 4\mathbf{I}) - 8(5\mathbf{A}^2 - 8\mathbf{A} + 4\mathbf{I}) + 4\mathbf{A}^2 \\
&= 25(5\mathbf{A}^2 - 8\mathbf{A} + 4\mathbf{I}) - 76\mathbf{A}^2 + 84\mathbf{A} - 32\mathbf{I} \\
&= 49\mathbf{A}^2 - 116\mathbf{A} + 68\mathbf{I}
\end{aligned}$$

Now calculate \mathbf{A}^2 and substitute. ■

Question 2(b) Let \mathbf{A} and \mathbf{B} be matrices of order n . If $\mathbf{I} - \mathbf{AB}$ is invertible, then $\mathbf{I} - \mathbf{BA}$ is also invertible and

$$(\mathbf{I} - \mathbf{BA})^{-1} = \mathbf{I} + \mathbf{B}(\mathbf{I} - \mathbf{AB})^{-1}\mathbf{A}$$

Show that \mathbf{AB} and \mathbf{BA} have the same characteristic values.

Solution.

$$\begin{aligned}
&(\mathbf{I} + \mathbf{B}(\mathbf{I} - \mathbf{AB})^{-1}\mathbf{A})(\mathbf{I} - \mathbf{BA}) \\
&= \mathbf{I} - \mathbf{BA} + \mathbf{B}(\mathbf{I} - \mathbf{AB})^{-1}\mathbf{A} - \mathbf{B}(\mathbf{I} - \mathbf{AB})^{-1}\mathbf{ABA} \\
&= [\mathbf{I} + \mathbf{B}(\mathbf{I} - \mathbf{AB})^{-1}\mathbf{A}] - \mathbf{B}[\mathbf{I} + (\mathbf{I} - \mathbf{AB})^{-1}\mathbf{AB}]\mathbf{A} \tag{1}
\end{aligned}$$

$$\text{Now } (\mathbf{I} - \mathbf{AB})^{-1}(\mathbf{I} - \mathbf{AB}) = (\mathbf{I} - \mathbf{AB})^{-1} - (\mathbf{I} - \mathbf{AB})^{-1}\mathbf{AB} = \mathbf{I}$$

$$\therefore (\mathbf{I} - \mathbf{AB})^{-1} = \mathbf{I} + (\mathbf{I} - \mathbf{AB})^{-1}\mathbf{AB}$$

$$\begin{aligned}
\text{Substituting in (1)} \quad &(\mathbf{I} + \mathbf{B}(\mathbf{I} - \mathbf{AB})^{-1}\mathbf{A})(\mathbf{I} - \mathbf{BA}) \\
&= \mathbf{I} + \mathbf{B}(\mathbf{I} - \mathbf{AB})^{-1}\mathbf{A} - \mathbf{B}(\mathbf{I} - \mathbf{AB})^{-1}\mathbf{A} = \mathbf{I}
\end{aligned}$$

Thus $\mathbf{I} - \mathbf{BA}$ is invertible and $(\mathbf{I} - \mathbf{BA})^{-1} = \mathbf{I} + \mathbf{B}(\mathbf{I} - \mathbf{AB})^{-1}\mathbf{A}$ as desired.

We shall show that $\lambda\mathbf{I} - \mathbf{AB}$ is invertible if and only if $\lambda\mathbf{I} - \mathbf{BA}$ is invertible. This means that if λ is an eigenvalue of \mathbf{AB} , then $|\lambda\mathbf{I} - \mathbf{AB}| = 0 \Rightarrow |\lambda\mathbf{I} - \mathbf{BA}| = 0$ so λ is an eigenvalue of \mathbf{BA} .

If $\lambda\mathbf{I} - \mathbf{AB}$ is invertible, then

$$\begin{aligned}
&(\mathbf{I} + \mathbf{B}(\lambda\mathbf{I} - \mathbf{AB})^{-1}\mathbf{A})(\lambda\mathbf{I} - \mathbf{BA}) \\
&= \lambda\mathbf{I} - \mathbf{BA} + \lambda\mathbf{B}(\lambda\mathbf{I} - \mathbf{AB})^{-1}\mathbf{A} - \mathbf{B}(\lambda\mathbf{I} - \mathbf{AB})^{-1}\mathbf{ABA} \\
&= \lambda[\mathbf{I} + \mathbf{B}(\lambda\mathbf{I} - \mathbf{AB})^{-1}\mathbf{A}] - \mathbf{B}[\mathbf{I} + (\lambda\mathbf{I} - \mathbf{AB})^{-1}\mathbf{AB}]\mathbf{A} \tag{2}
\end{aligned}$$

$$\text{Now } (\lambda\mathbf{I} - \mathbf{AB})^{-1}(\lambda\mathbf{I} - \mathbf{AB}) = \lambda(\lambda\mathbf{I} - \mathbf{AB})^{-1} - (\lambda\mathbf{I} - \mathbf{AB})^{-1}\mathbf{AB} = \mathbf{I}$$

$$\therefore \lambda(\lambda\mathbf{I} - \mathbf{AB})^{-1} = \mathbf{I} + (\lambda\mathbf{I} - \mathbf{AB})^{-1}\mathbf{AB}$$

$$\begin{aligned}
\text{Substituting in (2)} \quad &(\mathbf{I} + \mathbf{B}(\lambda\mathbf{I} - \mathbf{AB})^{-1}\mathbf{A})(\lambda\mathbf{I} - \mathbf{BA}) \\
&= \lambda\mathbf{I} + \lambda\mathbf{B}(\lambda\mathbf{I} - \mathbf{AB})^{-1}\mathbf{A} - \lambda\mathbf{B}(\lambda\mathbf{I} - \mathbf{AB})^{-1}\mathbf{A} = \lambda\mathbf{I}
\end{aligned}$$

Thus $\lambda\mathbf{I} - \mathbf{BA}$ is invertible if $\lambda\mathbf{I} - \mathbf{AB}$ is invertible. The converse is obvious as the situation is symmetric, thus \mathbf{AB} and \mathbf{BA} have the same eigenvalues.

We give another simple proof of the fact that \mathbf{AB} and \mathbf{BA} have the same eigenvalues.

1. Let 0 be an eigenvalue of \mathbf{AB} . This means that \mathbf{AB} is singular, i.e. $0 = |\mathbf{AB}| = |\mathbf{A}||\mathbf{B}| = |\mathbf{BA}|$, so \mathbf{BA} is singular, hence 0 is an eigenvalue of \mathbf{BA} .

2. Let $\lambda \neq 0$ be an eigenvalue of \mathbf{AB} and let $\mathbf{x} \neq \mathbf{0}$ be an eigenvector corresponding to λ , i.e. $\mathbf{ABx} = \lambda\mathbf{x}$. Let $\mathbf{y} = \mathbf{Bx}$. Then $\mathbf{y} \neq \mathbf{0}$, because $\mathbf{Ay} = \mathbf{ABx} = \lambda\mathbf{x} \neq \mathbf{0}$ as $\lambda \neq 0$. Now $\mathbf{BAy} = \mathbf{BABx} = \mathbf{B(ABx)} = \lambda\mathbf{Bx} = \lambda\mathbf{y}$. Thus λ is an eigenvalue of \mathbf{BA} . ■

Question 2(c) Let $a, b \in \mathbb{C}, |b| = 1$ and let \mathbf{H} be a Hermitian matrix. Show that the eigenvalues of $a\mathbf{I} + b\mathbf{H}$ lie on a straight line in the complex plane.

Solution. Let t be an eigenvalue of \mathbf{H} , which has to be real because \mathbf{H} is Hermitian. Clearly $a + tb$ is an eigenvalue of $a\mathbf{I} + b\mathbf{H}$. Conversely, if λ is an eigenvalue of $a\mathbf{I} + b\mathbf{H}$, then $\frac{\lambda - a}{b}$ (note $b \neq 0$ as $|b| = 1$) is an eigenvalue of \mathbf{H} .

Clearly $a + tb$ lies on the straight line joining points a and $a + b$:

$$z = (1 - x)a + x(b - a), \quad x \in \mathbb{R}$$

For the sake of completeness, we prove that the eigenvalues of a Hermitian matrix \mathbf{H} are real. Let $\mathbf{z} \neq \mathbf{0}$ be an eigenvector corresponding to the eigenvalue t .

$$\begin{aligned} \mathbf{Hz} &= t\mathbf{z} \\ \Rightarrow \bar{\mathbf{z}}'\mathbf{Hz} &= t\bar{\mathbf{z}}'\mathbf{z} \\ \Rightarrow \overline{\bar{\mathbf{z}}'\mathbf{Hz}} &= \overline{t\bar{\mathbf{z}}'\mathbf{z}} \\ \text{But } \overline{\bar{\mathbf{z}}'\mathbf{Hz}} &= \bar{\mathbf{z}}'\overline{\mathbf{H}\mathbf{z}} = \bar{\mathbf{z}}'\mathbf{Hz} = t\bar{\mathbf{z}}'\mathbf{z} \\ \Rightarrow t\bar{\mathbf{z}}'\mathbf{z} &= \bar{t}\bar{\mathbf{z}}'\mathbf{z} \\ \Rightarrow t &= \bar{t} \quad \because \bar{\mathbf{z}}'\mathbf{z} \neq 0 \end{aligned}$$

Question 3(a) Let \mathbf{A} be a symmetric matrix. Show that \mathbf{A} is positive definite if and only if its eigenvalues are all positive.

Solution. \mathbf{A} is real symmetric so all eigenvalues of \mathbf{A} are real. Let $\lambda_1, \lambda_2, \dots, \lambda_n$ be eigenvalues of \mathbf{A} , not necessarily distinct. Let \mathbf{x}_1 be an eigenvector corresponding to λ_1 . Since λ_1 and \mathbf{A} are real, \mathbf{x}_1 is also real. Replacing \mathbf{x}_1 if necessary by $\mu\mathbf{x}_1$, μ suitable, we can assume that $\|\mathbf{x}_1\| = \sqrt{\mathbf{x}_1'\mathbf{x}_1} = 1$.

Let \mathbf{P}_1 be an orthogonal matrix with \mathbf{x}_1 as its first column. Such a \mathbf{P}_1 exists, as will be shown at the end of this result. Clearly the first column of the matrix $\mathbf{P}_1^{-1}\mathbf{AP}_1$ is equal to $\mathbf{P}_1^{-1}\mathbf{Ax} = \lambda_1\mathbf{P}_1^{-1}\mathbf{x} = \begin{pmatrix} \lambda_1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$, because $\mathbf{P}_1^{-1}\mathbf{x}$ is the first column of $\mathbf{P}_1^{-1}\mathbf{P} = \mathbf{I}$. Thus $\mathbf{P}_1^{-1}\mathbf{AP}_1 = \begin{pmatrix} \lambda_1 & \mathbf{L} \\ \mathbf{0} & \mathbf{B} \end{pmatrix} = \mathbf{P}'_1\mathbf{AP}_1$ where \mathbf{B} is $(n - 1) \times (n - 1)$ symmetric. Since $\mathbf{P}'_1\mathbf{AP}_1$ is symmetric, it follows that $\mathbf{P}_1^{-1}\mathbf{AP}_1 = \mathbf{P}'_1\mathbf{AP}_1 = \begin{pmatrix} \lambda_1 & \mathbf{0} \\ \mathbf{0} & \mathbf{B} \end{pmatrix}$. Induction now gives that there

exists an $(n - 1) \times (n - 1)$ orthogonal matrix \mathbf{Q} such that $\mathbf{Q}'\mathbf{BQ} = \begin{pmatrix} \lambda_2 & 0 & \dots & 0 \\ \dots & & & \\ 0 & 0 & \dots & \lambda_n \end{pmatrix}$

where $\lambda_2, \lambda_3, \dots, \lambda_n$ are eigenvalues of \mathbf{B} . Let $\mathbf{P}_2 = \begin{pmatrix} 1 & \mathbf{0} \\ \mathbf{0} & \mathbf{Q} \end{pmatrix}$, then \mathbf{P}_2 is orthogonal and $\mathbf{P}_2' \mathbf{P}_1' \mathbf{A} \mathbf{P}_1 \mathbf{P}_2 = \text{diagonal}[\lambda_1, \dots, \lambda_n]$. Set $\mathbf{P} = \mathbf{P}_1 \mathbf{P}_2 \dots \mathbf{P}_n$, and $(y_1, \dots, y_n) \mathbf{P}' = \mathbf{x}$ then $\mathbf{x}' \mathbf{A} \mathbf{x} = \mathbf{y}' \mathbf{P}' \mathbf{A} \mathbf{P} \mathbf{y} = \sum_{i=1}^n \lambda_i^2 y_i^2$.

Since \mathbf{P} is non-singular, quadratic forms $\mathbf{x}' \mathbf{A} \mathbf{x}$ and $\sum_{i=1}^n \lambda_i^2 y_i^2$ assume the same values. Hence \mathbf{A} is positive definite if and only if $\sum_{i=1}^n \lambda_i^2 y_i^2$ is positive definite if and only if $\lambda_i > 0$ for all i .

Result used: If \mathbf{x}_1 is a real vector such that $\|\mathbf{x}_1\| = \sqrt{\mathbf{x}_1' \mathbf{x}_1} = 1$ then there exists an orthogonal matrix with \mathbf{x}_1 as its first column.

Proof: We have to find real column vectors $\mathbf{x}_2, \dots, \mathbf{x}_n$ such that $\|\mathbf{x}_i\| = 1, 2 \leq i \leq n$ and $\mathbf{x}_2, \dots, \mathbf{x}_n$ is an orthonormal system i.e. $\mathbf{x}_i' \mathbf{x}_j = 0, i \neq j$. Consider the single equation $\mathbf{x}_1' \mathbf{x} = 0$, where \mathbf{x} is a column vector to be determined. This equation has a non-zero solution, in fact the space of solutions is of dimension $n - 1$, the rank of the coefficient matrix being 1. If \mathbf{y}_2 is a solution, we take $\mathbf{x}_2 = \frac{\mathbf{y}_2}{\|\mathbf{y}_2\|}$ so that $\mathbf{x}_1' \mathbf{x}_2 = 0$.

We now consider the two equations $\mathbf{x}_1' \mathbf{x} = 0, \mathbf{x}_2' \mathbf{x} = 0$. Again the number of unknowns is more than the number of equations, so there is a solution, say \mathbf{y}_3 , and take $\mathbf{x}_3 = \frac{\mathbf{y}_3}{\|\mathbf{y}_3\|}$ to get $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3$ mutually orthogonal.

Proceeding in this manner, if we consider $n - 1$ equations $\mathbf{x}_1' \mathbf{x} = 0, \dots, \mathbf{x}_{n-1}' \mathbf{x} = 0$, these will have a nonzero solution \mathbf{y}_n , so we set $\mathbf{x}_n = \frac{\mathbf{y}_n}{\|\mathbf{y}_n\|}$. Clearly $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ is an orthonormal system, and therefore $\mathbf{P} = [\mathbf{x}_1, \dots, \mathbf{x}_n]$ is an orthogonal matrix having \mathbf{x}_1 as a first column. ■

Question 3(b) Let \mathbf{A} and \mathbf{B} be square matrices of order n , show that $\mathbf{AB} - \mathbf{BA}$ can never be equal to the identity matrix.

Solution. Let $\mathbf{A} = \langle a_{ij} \rangle$ and $\mathbf{B} = \langle b_{ij} \rangle$. Then

$$\begin{aligned} \text{tr } \mathbf{AB} &= \text{Sum of diagonal elements of } \mathbf{AB} \\ &= \sum_{i=1}^n \sum_{k=1}^n a_{ik} b_{ki} = \sum_{k=1}^n \sum_{i=1}^n b_{ki} a_{ik} = \text{tr } \mathbf{BA} \end{aligned}$$

Thus $\text{tr}(\mathbf{AB} - \mathbf{BA}) = \text{tr } \mathbf{AB} - \text{tr } \mathbf{BA} = 0$. But the trace of the identity matrix is n , thus $\mathbf{AB} - \mathbf{BA}$ can never be equal to the identity matrix. ■

Question 3(c) Let $\mathbf{A} = \langle a_{ij} \rangle, 1 \leq i, j \leq n$. If $\sum_{\substack{j=1 \\ i \neq j}}^n |a_{ij}| < |a_{ii}|$, then the eigenvalues of \mathbf{A} lie in the disc

$$|\lambda - a_{ii}| \leq \sum_{\substack{j=1 \\ i \neq j}}^n |a_{ij}|$$

Solution. See the solution to question 2(c), year 1997. We showed that if $|\lambda - a_{ii}| > \sum_{\substack{j=1 \\ i \neq j}}^n |a_{ij}|$ then $|\lambda \mathbf{I} - \mathbf{A}| \neq 0$, so λ is not an eigenvalue of \mathbf{A} . Thus if λ is an eigenvalue, then $|\lambda - a_{ii}| \leq \sum_{\substack{j=1 \\ i \neq j}}^n |a_{ij}|$, so λ lies in the disc described in the question. ■